



ARDB 2019. AFRICAN RAPTOR DATABANK:
a secure, live data observatory for the
distribution and movements of African raptors.
Habitat Info Ltd, Solva, UK.
www.habitatinfo.com/ardb_observatory/
02/12/2019: 16:10:07

1200Pel's Fishing-owl

Scotopelia peli

IUCN Status Least Concern
IUCN (2019-2)

male mass = g
female mass = 3000 g
mean mass = g

area requirement (of pair/2.2 individuals) = 60.4 km²

from Newton (1979) figure 10: $area\ km^2 = 0.4 + (0.02 * female\ mass\ g)$
replaced by direct measurement of field density where we have data



this figure is used for estimating density of breeding individuals and offspring in good habitat
we assume twice the area requirement in suboptimal habitat
and half the area requirement in optimal habitat
breeding population estimates represent theoretical maxima where habitat is saturated
nest-site limitation and the floating / non-breeding population are not yet fully accounted for

HABITAT SPACE ANALYSIS

	extent (km ²)	individuals	pairs
suboptimal 25-50% suitable	3825439		
good habitat 50-70% suitable	908432		
optimal habitat 70-100% suitable	286201		
total habitat 25-100% suitable	5020072		

reporting rate from mobile app survey data

2014 (26724 km) : birds per 100 km
2015 (87188 km) : birds per 100 km
2016 (81150 km) : birds per 100 km
2017 (57721 km) : birds per 100 km
2018 (19798 km) : birds per 100 km



Legend

- × vagrant records (0)
- recent observations (182)
- ★ historic breeding records (0)
- movement data (0)
- historic sightings / specimens (136)
- large protected areas

range_type

- ▨ undefined
- historic occurrence
- breeding only, sparse occurrence
- breeding season only
- non-breeding only, sparse occurrence
- non-breeding season only
- only on migration / vagrants
- resident
- resident, sparse occurrence

habitat suitability

- Value
- 100 % suitable
 - 0 % suitable

0 550 1,100 2,200 Kilometers

Lambert Azimuth Equal Area Projection (Central Meridian 20)