



ARDB 2019. AFRICAN RAPTOR DATABANK:
a secure, live data observatory for the
distribution and movements of African raptors.
Habitat Info Ltd, Solva, UK.
www.habitatinfo.com/ardb_observatory/
02/12/2019: 13:00:05

430 Eurasian Buzzard

Buteo buteo

IUCN Status Least Concern
IUCN (2019-2)

male mass = 800 g
female mass = 950 g
mean mass = 950 g

area requirement (of pair/2.2 individuals) = 19.4 km²

from Newton (1979) figure 10: $area\ km^2 = 0.4 + (0.02 * female\ mass\ g)$
replaced by direct measurement of field density where we have data



this figure is used for estimating density of breeding individuals and offspring in good habitat
we assume twice the area requirement in suboptimal habitat
and half the area requirement in optimal habitat
breeding population estimates represent theoretical maxima where habitat is saturated
nest-site limitation and the floating / non-breeding population are not yet fully accounted for

HABITAT SPACE ANALYSIS

	extent (km ²)	individuals	pairs
suboptimal 25-50% suitable	476227	27003	0
good habitat 50-70% suitable	195267	22144	0
optimal habitat 70-100% suitable	16264	3689	0
total habitat 25-100% suitable	687758	52835	0

reporting rate from mobile app survey data

2014 (26724 km) : birds per 100 km
2015 (87188 km) : 0.001147 birds per 100 km
2016 (81150 km) : 0.045595 birds per 100 km
2017 (57721 km) : 0.069299 birds per 100 km
2018 (19798 km) : 0.171739 birds per 100 km



Legend

- × vagrant records (0)
- recent observations (536)
- ★ historic breeding records (0)
- movement data (0)
- historic sightings / specimens (0)
- large protected areas

range_type

- undefined
- historic occurrence
- breeding only, sparse occurrence
- breeding season only
- non-breeding only, sparse occurrence
- non-breeding season only
- only on migration / vagrants
- resident
- resident, sparse occurrence

habitat suitability

Value

- 100 % suitable
- 0 % suitable